

Research Article

A relation between Wiener index and Mostar index for daisy cubes

Michel Mollard*

Institut Fourier, CNRS, Université Grenoble Alpes, Grenoble, France

(Received: 12 May 2022. Received in revised form: 15 June 2022. Accepted: 16 June 2022. Published online: 18 June 2022.)

© 2022 the author. This is an open access article under the CC BY (International 4.0) license (www.creativecommons.org/licenses/by/4.0/).

Abstract

Daisy cubes form a class of isometric subgraphs of the hypercubes Q_n . They include some previously well-known families of graphs like Fibonacci cubes and Lucas cubes. Daisy cubes also appear in chemical graph theory. Two distance invariants, Wiener and Mostar indices, have been introduced in the context of the mathematical chemistry. The Wiener index $W(G)$ is the sum of distance between all unordered pairs of vertices of a graph G . The Mostar index $Mo(G)$ is a measure of how far G is from being distance balanced. In this paper, it is proved that the Wiener and Mostar indices of a daisy cube G are linked by the relation $2W(G) - Mo(G) = |V(G)| \cdot |E(G)|$. An expression concerning the Wiener and Mostar indices for daisy cubes is also deduced.

Keywords: daisy cube; Wiener index; Mostar index; Fibonacci cube; partial cube.

2020 Mathematics Subject Classification: 05C09, 05C12, 05C92.

1. Introduction

The *Fibonacci cube* of dimension n , denoted as Γ_n , is the subgraph of the hypercube Q_n induced by vertices with no consecutive 1's. This graph was introduced in [6] as an interconnection network. Lucas cubes (see [17]) are the cyclic version of Fibonacci cubes. Structural properties of these graphs have been widely studied (see [10] for a survey). Fibonacci cubes also play a role in mathematical chemistry. Indeed they are precisely the resonance graphs of fibonacenes, an important class of hexagonal chains [15]. Later in [21], a similar connection have been found between Lucas cubes and the resonance graphs of cyclic polyphenanthrenes, which are related to non-cyclic fibonacenes. Fibonacci cubes and Lucas cubes belong to daisy cubes [13], a family of isometric subgraphs of hypercubes. The connection between daisy cubes and resonance graphs of catacondensed even ring systems have been explored in [2].

The Wiener index $W(G)$ of a connected graph G is the sum of distance of all unordered pairs of vertices of G . This distance invariant is important in mathematical chemistry. The Wiener index of Γ_n and Λ_n have been determined in [12]. Recently the Mostar index $Mo(G)$ has been introduced in [3] again in the context of graph chemical theory. It measures how far G is from being distance-balanced. The Mostar index of Fibonacci and Lucas cubes have been determined in [4]. See [1] for a recent survey on Mostar index. In this note we prove that if G is a daisy cube, then the Wiener and the Mostar index are linked by the relation

$$2W(G) - Mo(G) = |V(G)| \cdot |E(G)|.$$

In the last section we derive similar expressions for $W(G)$ and $Mo(G)$ from the sequence of the number of edges using the direction i for $i \in [n]$.

2. Preliminaries

In this section, some concepts and notations needed in the rest of this paper are given. We denote by $[n]$ the set of integers i such that $1 \leq i \leq n$. Let $\{F_n\}$ be the *Fibonacci numbers*: $F_0 = 0$, $F_1 = 1$, $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$. Let $B = \{0, 1\}$. It will be convenient to identify elements $u = (u_1, \dots, u_n) \in B^n$ and strings of length n over B . We thus briefly write u as $u_1 \dots u_n$ and call u_i the i th coordinate of u . We will use the power notation for the concatenation of bits, for instance $0^n = 0 \dots 0 \in B^n$. We will denote by \bar{u}_i the binary complement of u_i .

The vertex set of Q_n , the *hypercube of dimension n* , is the set B^n , two vertices being adjacent if and only if they differ in precisely one coordinate. We will say that an edge uv of Q_n uses the direction i if u and v differ in the coordinate i .

*E-mail address: michel.mollard@univ-grenoble-alpes.fr

The *distance* between two vertices u and v of a graph G is the number of edges on a shortest shortest u, v -path. It is immediate that the distance between two vertices of Q_n is the number of coordinates the strings differ, sometime called Hamming distance. The *Wiener index* $W(G)$ of a connected graph G is defined as the sum of all distances between pairs of vertices of G . Hence,

$$W(G) = \sum_{\{u,v\} \subseteq V(G)} d(u,v).$$

A *Fibonacci string* of length n is a binary string $b = b_1b_2 \dots b_n$ with $b_i \cdot b_{i+1} = 0$ for $1 \leq i < n$. In other words a Fibonacci string is a binary string without 11 as substring. The *Fibonacci cube* Γ_n ($n \geq 1$) is the subgraph of Q_n induced by \mathcal{F}_n the set of Fibonacci strings of length n . Because of the empty string ϵ , $\Gamma_0 = K_1$. Note that for any integer n , $|V(\Gamma_n)| = F_{n+2}$. A Fibonacci string b of length n is a *Lucas string* if $b_1 \cdot b_n \neq 1$. That is, a Lucas string has no two consecutive 1s including the first and the last elements of the string. The *Lucas cube* Λ_n is the subgraph of Q_n induced by the Lucas strings of length n . We have $\Lambda_0 = \Lambda_1 = K_1$.

Fibonacci cubes and Lucas cubes were extended to generalized Fibonacci cubes [8] and generalized Lucas cubes [7]. For any arbitrary string f the generalized Fibonacci cubes $Q_n[f]$ is the subgraph of Q_n induced by strings of B^n which do not contain f as substring. Similarly the generalized Lucas cubes $Q_n[\overleftarrow{f}]$ is induced by strings without a circulation containing f as substring. Classical Fibonacci and Lucas cubes correspond to $f = 11$.

If u and v are vertices of a graph G , the *interval* $I_G(u, v)$ between u and v (in G) is the set of vertices lying on shortest u, v -path, that is, $I_G(u, v) = \{w | d(u, v) = d(u, w) + d(w, v)\}$. We will also write $I(u, v)$ when G will be clear from the context. A subgraph G of a graph H is an *isometric subgraph* if the distance between any vertices of G equals the distance between the same vertices in H . Isometric subgraphs of hypercubes are called *partial cubes*. The *dimension* of a partial cube G is the smallest integer d such that G is an isometric subgraph of Q_d . Many important classes of graphs are partial cubes, in particular trees, median graphs, benzenoid graphs, phenylenes, grid graphs and bipartite torus graphs. In addition, Fibonacci and Lucas cubes are partial cubes as well, see [9].

If $G = (V(G), E(G))$ is a graph and $X \subseteq V(G)$, then $\langle X \rangle$ denotes the subgraph of G induced by X . Let \leq be a partial order on B^n defined with $u_1 \dots u_n \leq v_1 \dots v_n$ if $u_i \leq v_i$ holds for $i \in [n]$. For $X \subseteq B^n$ we define the graph $Q_n(X)$ as the subgraph of Q_n with

$$Q_n(X) = \langle \{u \in B^n | u \leq x \text{ for some } x \in X\} \rangle$$

and say that $Q_n(X)$ is the *daisy cube generated by X* . Note that if \widehat{X} is the antichain consisting of the maximal elements of the poset (X, \leq) , then $Q_n(\widehat{X}) = Q_n(X)$. As noticed in the daisy cube introductory paper [13] we can alternatively say that

$$Q_n(X) = \left\langle \bigcup_{x \in X} I_{Q_n}(x, 0^n) \right\rangle = \left\langle \bigcup_{x \in \widehat{X}} I_{Q_n}(x, 0^n) \right\rangle.$$

Finally we will say that a graph G is a *daisy cube* if there exist an isometric embedding of G in some hypercube Q_n and a subset X of B^n such that G is the daisy cube generated by X . Such an embedding will be called a *proper embedding*.

By construction daisy cubes are partial cubes and as noticed in the same paper Fibonacci cubes, Lucas cubes, bipartite wheels, vertex-deleted cubes and hypercubes themselves are daisy cubes. It is easy to see that Pell graphs [16] are also daisy cubes. Furthermore A. Vesel proved [19] that the cube complement of a daisy cube is a daisy cube.

For any fixed integer $s \geq 2$ the generalized Fibonacci cubes $Q_n[1^s]$ and Lucas cubes $Q_n[\overleftarrow{1^s}]$ are also daisy cubes. The Wiener index of these two families of graphs have been studied in [14]. A construction of daisy cubes in terms of expansion procedure is given in [18].

Let a partial cube G of dimension n be given together with its isometric embedding into Q_n . Then for $i = 1, 2, \dots, n$ and $\chi = 0, 1$ the *semicube* $W_{(i,\chi)}$ is defined as follows:

$$W_{(i,\chi)}(G) = \{u = u_1u_2 \dots u_n \in V(G) \mid u_i = \chi\}.$$

For a fixed i , the pair $W_{(i,0)}(G), W_{(i,1)}(G)$ of semicubes is called a *complementary pair of semicubes*. The Wiener index of partial cubes can be determined using the following result [11].

Theorem 2.1. *Let G be a partial cube of dimension n isometrically embedded into Q_n . Then*

$$W(G) = \sum_{i=1}^n |W_{(i,0)}(G)| \cdot |W_{(i,1)}(G)|.$$

Notice that, as expected, this expression is independent of the embedding. Indeed isometric embeddings of G into Q_n are unique up to the automorphisms of Q_n , see [20]. These automorphisms are generated by $\Theta = \{\theta_{i,j} | i, j \in [n]\}$ and

$T = \{\tau_i | i \in [n]\}$ where $\theta_{i,j}$ is the transposition of coordinates i and j and τ_i is the complementation of the coordinate i . A permutation $\theta_{i,j}$ induces a similar permutation on complementary pairs and an element of T is also without effect on the expression of $W(G)$ since τ_i exchanges $W_{(i,0)}(G)$ and $W_{(i,1)}(G)$.

Let $G = (V(G), E(G))$ be a connected graph. For any edge $uv \in E(G)$, let $n_{u,v}$ denote the number of vertices which are closer to u than to v . The *Mostar index* $Mo(G)$ is defined as

$$Mo(G) = \sum_{uv \in E(G)} |n_{u,v} - n_{v,u}|.$$

An expression of the Mostar index for partial cubes is given in [5].

3. A relation

This section is devoted to the proof of the following theorem.

Theorem 3.1. *Let G be a daisy cube. Then the Wiener and the Mostar indices of G are linked by the relation*

$$2W(G) - Mo(G) = |V(G)| \cdot |E(G)|.$$

Like in [12] where it is applied in particular to Fibonacci cubes and Lucas cubes, Theorem 2.1 will be our starting point for proving the relation. Consider a proper embedding of G in Q_n and thus let $X \subseteq B^n$ such that

$$G = \langle \{u \in B^n | u \leq x \text{ for some } x \in X\} \rangle.$$

For any $i \in [n]$ let E_i be the set of edges of G using the direction i .

Proposition 3.1. *For any $i \in [n]$ we have*

$$|W_{(i,0)}(G)| \geq |W_{(i,1)}(G)|.$$

Proof. Let $u = u_1u_2 \dots u_n$ in $W_{(i,1)}(G)$ and consider $\theta(u) = u_1 \dots u_{i-1}0u_{i+1} \dots u_n$. Note that $\theta(u) \leq u$ and since u is a vertex of G there exists $x \in X$ with $u \leq x$. Therefore $\theta(u) \leq x$ and $\theta(u) \in V(G)$. By this way we construct an injective mapping from $W_{(i,1)}(G)$ to $W_{(i,0)}(G)$. □

Proposition 3.2. *For any $i \in [n]$ we have*

$$|E_i| = |W_{(i,1)}(G)|.$$

Proof. Indeed let $u = u_1u_2 \dots u_n$ in $W_{(i,1)}(G)$ and let $v = u_1 \dots u_{i-1}0u_{i+1} \dots u_n$. Using the same argument as in the previous proof, it is clear that v is a vertex of G and that the edge uv belongs to E_i . Reciprocally exactly one of the extremities of a given edge of E_i belongs to $W_{(i,1)}(G)$. We obtain a bijective mapping between $W_{(i,1)}(G)$ and E_i . □

Proposition 3.3. *For any edge uv of E_i with $u_i = 0$ we have*

$$W_{(i,0)}(G) = \{w \in V(G) | d(w, u) < d(w, v)\},$$

$$W_{(i,1)}(G) = \{w \in V(G) | d(w, v) < d(w, u)\}.$$

Proof. Since $d(w, u)$ and $d(w, v)$ are the number of coordinates the strings differ and since u, v differ only by the coordinate i it is clear that $d(w, u) = d(w, v) + 1$ if $w \in W_{(i,1)}(G)$ and $d(w, u) = d(w, v) - 1$ otherwise. □

Lemma 3.1. *Let G be a daisy cube of dimension n properly embedded into Q_n . Then*

$$Mo(G) = \sum_{i=1}^n |W_{(i,1)}(G)| (|W_{(i,0)}(G)| - |W_{(i,1)}(G)|).$$

Proof. Let $e = uv$ be an edge of E_i with $u_i = 0$. By Propositions 3.3 and 3.1 we have $n_{u,v} = |W_{(i,0)}(G)| \geq |W_{(i,1)}(G)| = n_{v,u}$. The contribution of the edge e to $\sum_{uv \in E(G)} |n_{u,v} - n_{v,u}|$ is thus $|W_{(i,0)}(G)| - |W_{(i,1)}(G)|$. Therefore

$$Mo(G) = \sum_{i=1}^n \sum_{uv \in E_i} (|W_{(i,0)}(G)| - |W_{(i,1)}(G)|) = \sum_{i=1}^n |E_i| (|W_{(i,0)}(G)| - |W_{(i,1)}(G)|).$$

The conclusion follows from Proposition 3.2. □

We can now proceed to the proof of our relation given in Theorem 3.1. By Theorem 2.1 we obtain

$$W(G) = \sum_{i=1}^n |W_{(i,0)}(G)| \cdot |W_{(i,1)}(G)|.$$

Since $|W_{(i,0)}(G)| + |W_{(i,1)}(G)| = |V(G)|$ we deduce from Lemma 3.1 that

$$\begin{aligned} 2W(G) - Mo(G) &= \sum_{i=1}^n |W_{(i,1)}(G)| (2|W_{(i,0)}(G)| - |W_{(i,0)}(G)| + |W_{(i,1)}(G)|) \\ &= \sum_{i=1}^n |W_{(i,1)}(G)| \cdot |V(G)|. \end{aligned}$$

Since $E(G)$ is the disjoint union $E(G) = \cup_{i=1}^n E_i$ we deduce from Proposition 3.2 that $\sum_{i=1}^n |W_{(i,1)}(G)| = |E(G)|$ and the relation follows. \square

4. Conclusion

The Wiener and Mostar indices of daisy cubes are completely determined by $|V(G)|$ and the sequence $|E_i|$ (for $i \in [n]$) of the number of edges using the direction i which is identical to the sequence of $|W_{(i,1)}(G)|$. Indeed, from Theorem 2.1 and Lemma 3.1 we have the relation

$$W(G) - Mo(G) = \sum_{i=1}^n |W_{(i,1)}(G)|^2.$$

Combining this identity with that of Theorem 3.1 we obtain the following assertion.

Corollary 4.1. *Let G be a daisy cube properly embedded into Q_n . For $i \in [n]$ let E_i be the set of edges using the direction i . Then the Wiener and the Mostar indices of G are*

$$\begin{aligned} W(G) &= |V(G)| \cdot |E(G)| - \sum_{i=1}^n |E_i|^2, \\ Mo(G) &= |V(G)| \cdot |E(G)| - 2 \sum_{i=1}^n |E_i|^2. \end{aligned}$$

In conclusion of this paper note that it will be interesting to give bijective proofs of Theorem 3.1 and Corollary 4.1.

References

- [1] A. Ali, T. Došlić, Mostar index: results and perspectives, *Appl. Math. Comput.* **404** (2021) #126245.
- [2] S. Brezovnik, N. Tratnik, P. Žigert, Resonance graphs of catacondensed even ring systems, *Appl. Math. Comput.* **374** (2020) #125064.
- [3] T. Došlić, I. Martinjak, R. Škrekovski, S. T. Spužević, I. Zubac, Mostar Index, *J. Math. Chem.* **56** (2018) 2995–3013.
- [4] Ö. Egecioğlu, E. Saygi, Z. Saygi, The Mostar index of Fibonacci and Lucas cubes, *Bull. Malays. Math. Sci. Soc.* **44** (2021) 3677–3687.
- [5] M. Ghorbani, S. Rahmani, M. Eslampoor, Some new results on Mostar index of graphs, *Iranian J. Math. Chem.* **11** (2020) 33–42.
- [6] W. J. Hsu, Fibonacci cubes a new interconnection technology, *IEEE Trans. Parallel Distrib. Syst.* **4** (1993) 3–12.
- [7] A. Ilić, S. Klavžar, Y. Rho, Generalized Lucas cubes, *Appl. Anal. Discrete Math.* **6** (2012) 82–94.
- [8] A. Ilić, S. Klavžar, Y. Rho, Generalized Fibonacci cubes, *Discrete Math.* **312** (2012) 2–11.
- [9] S. Klavžar, On median nature and enumerative properties of Fibonacci-like cubes, *Discrete Math.* **299** (2005) 145–153.
- [10] S. Klavžar, Structure of Fibonacci cubes: a survey, *J. Comb. Optim.* **25** (2013) 505–522.
- [11] S. Klavžar, I. Gutman, B. Mohar, Labeling of benzenoid systems which reflects the vertex-distance relations, *J. Chem. Inf. Comput. Sci.* **35** (1995) 590–593.
- [12] S. Klavžar, M. Mollard, Wiener index and Hosoya polynomials of Fibonacci and Lucas cubes, *MATCH Commun. Math. Comput. Chem.* **68** (2012) 311–324.
- [13] S. Klavžar, M. Mollard, Daisy cubes and distance polynomial, *European J. Combin.* **80** (2019) 214–223.
- [14] S. Klavžar, Y. Rho, On the Wiener index of generalized Fibonacci and Lucas cubes, *Discrete Appl. Math.* **187** (2015) 155–160.
- [15] S. Klavžar, P. Žigert, Fibonacci cubes are the resonance graphs of fibonaccenes, *Fibonacci Quart.* **43** (2005) 269–276.
- [16] E. Munarini, Pell graphs, *Discrete Math.* **342** (2019) 2415–2428.
- [17] E. Munarini, C. P. Cippo, N. Z. Salvi, On the Lucas cubes, *Fibonacci Quart.* **39** (2001) 12–21.
- [18] A. Taranenko, Daisy cubes: A characterization and a generalization, *European J. Combin.* **85** (2020) #103058.
- [19] A. Vesel, Cube-complement of generalized Fibonacci cubes, *Discrete Math.* **342** (2019) 1139–1146.
- [20] P. M. Winkler, Isometric embedding in products of complete graphs, *Discrete Appl. Math.* **7** (1984) 221–225.
- [21] P. Žigert, M. Berlić, Lucas cubes and resonance graphs of cyclic polyphenantrenes, *MATCH Commun. Math. Comput. Chem.* **68** (2012) 77–90.